

## EDUVISTA MASTER INTERACTIVE INFOGRAPHIC &amp; PYQ COMPENDIUM

## CLASS 11 NCERT CHEMISTRY — CHAPTER 7: REDOX REACTIONS

Academic Standard: CBSE Board Exams + JEE Main + NEET Competitive Level

Official Branding: EDUVISTA | Learn • Understand • Achieve

OFFICIAL EDUVISTA INTERACTIVE LEARNING SYSTEM

## EDUVISTA INTERACTIVE SMART NOTES: REDOX REACTIONS

Class 11 NCERT Chemistry | Code: **kech201** | 100% Concept & Exam Mastery

✓ Interactive Decision Trees   ✓ Self-Testing PYQ Practice   ✓ Hyperlinked Jump Grid

✓ Step-by-Step Balancing Workbench

## INTERACTIVE JUMP MATRIX — QUICK SECTION NAVIGATOR

1. Classical &amp; Electronic Concepts

2. Competitive Electron Transfer

3. Oxidation Number Rules

4. Structural Paradoxes ( $\text{CrO}_5$ ,  $\text{H}_2\text{SO}_5$ )

5. Types of Redox &amp; Disproportionation

6. Balancing Workbench (Half-Reaction)

7. Redox Titrations &amp; Indicators

8. Galvanic Cell &amp; Daniell Cell

9. Electrochemical Series &  $E^\circ_{\text{cell}}$ 

10. CBSE Board Solved PYQs

11. JEE Main &amp; NEET Solved PYQs

12. Interactive Self-Test Quiz

## 1. CLASSICAL &amp; ELECTRONIC CONCEPTS OF REDOX

Redox reactions (Reduction-Oxidation) involve simultaneous electron transfer between reacting chemical species.

**Oxidation (OIL = Oxidation Is Loss of Electrons)**

- **Classical View:** Addition of Oxygen or Electronegative element; Removal of Hydrogen or Electropositive element.
- **Electronic View:** Loss of one or more electrons (Increases Oxidation Number).
- **Example:**  $2\text{Mg(s)} + \text{O}_2\text{(g)} \rightarrow 2\text{MgO(s)}$  |  $\text{Mg} \rightarrow \text{Mg}^{2+} + 2\text{e}^-$

**Reduction (RIG = Reduction Is Gain of Electrons)**

- **Classical View:** Removal of Oxygen or Electronegative element; Addition of Hydrogen or Electropositive element.
- **Electronic View:** Gain of one or more electrons (Decreases Oxidation Number).
- **Example:**  $2\text{HgO(s)} \rightarrow 2\text{Hg(l)} + \text{O}_2\text{(g)}$  ( $\Delta$ ) |  $\text{Hg}^{2+} + 2\text{e}^- \rightarrow \text{Hg}$

**Oxidising Agent vs Reducing Agent**

- **Oxidising Agent (Oxidant):** Accepts electrons → Gets reduced itself → Oxidation state decreases.
- **Reducing Agent (Reductant):** Donates electrons → Gets oxidised itself → Oxidation state increases.

[Back to Interactive Index](#)**2. COMPETITIVE ELECTRON TRANSFER & REACTIVITY SERIES**

Placing metallic strips in aqueous salt solutions demonstrates relative electron-releasing tendencies.

| COMPETITIVE ELECTRON TRANSFER EXPERIMENTS |                                                                                                          |                              |  |
|-------------------------------------------|----------------------------------------------------------------------------------------------------------|------------------------------|--|
| 1. Zn in CuSO <sub>4</sub> sol            | : $\text{Zn(s)} + \text{Cu}^{2+}\text{(aq)} \rightarrow \text{Zn}^{2+}\text{(aq)} + \text{Cu(s)}$        | [Zn loses $\text{e}^-$ > Cu] |  |
| 2. Cu in AgNO <sub>3</sub> sol            | : $\text{Cu(s)} + 2\text{Ag}^+\text{(aq)} \rightarrow \text{Cu}^{2+}\text{(aq)} + 2\text{Ag(s)}$         | [Cu loses $\text{e}^-$ > Ag] |  |
| 3. Co in NiSO <sub>4</sub> sol            | : $\text{Co(s)} + \text{Ni}^{2+}\text{(aq)} \rightleftharpoons \text{Co}^{2+}\text{(aq)} + \text{Ni(s)}$ | [Equilibrium]                |  |

**EDUVISTA MEMORY TRICK (ELECTRON RELEASE POWER):**

**Zn > Cu > Ag** (Zinc displaces Copper; Copper displaces Silver)

[Back to Interactive Index](#)**3. OXIDATION NUMBER (O.N.) DECISION TREE & RULES****INTERACTIVE DECISION TREE: HOW TO ASSIGN OXIDATION NUMBERS**

1. **Is the element uncombined/free state?** → O.N. = 0 (e.g. O<sub>2</sub>, S<sub>8</sub>, P<sub>4</sub>, Na, Fe).
2. **Is it a monoatomic ion?** → O.N. = Net Charge (e.g. Na<sup>+</sup> = +1, Mg<sup>2+</sup> = +2, Cl<sup>-</sup> = -1).
3. **Is the atom Fluorine?** → O.N. = -1 ALWAYS (No exceptions!).
4. **Is the atom Oxygen?**
  - In usual oxides (H<sub>2</sub>O, CO<sub>2</sub>) → O.N. = -2
  - In Peroxides (H<sub>2</sub>O<sub>2</sub>, Na<sub>2</sub>O<sub>2</sub>) → O.N. = -1
  - In Superoxides (KO<sub>2</sub>) → O.N. = -1/2
  - In Fluorides (OF<sub>2</sub>, O<sub>2</sub>F<sub>2</sub>) → O.N. = +2, +1

**5. Is the atom Hydrogen?**

- Bonded to Non-Metals (HCl, H<sub>2</sub>O) → O.N. = +1
- Bonded to Metals / Hydrides (LiH, NaH, CaH<sub>2</sub>) → O.N. = -1

| Rule #        | Element / Group            | Standard Oxidation Number    | Exceptions & Special Cases                                                                                                                                                      |
|---------------|----------------------------|------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Rule 1</b> | Free / Uncombined Elements | <b>Always 0</b>              | H <sub>2</sub> , O <sub>2</sub> , N <sub>2</sub> , S <sub>8</sub> , P <sub>4</sub> , Na, Fe (O.N. = 0)                                                                          |
| <b>Rule 2</b> | Monoatomic Ions            | <b>Equal to Ion Charge</b>   | Na <sup>+</sup> (+1), Mg <sup>2+</sup> (+2), Fe <sup>3+</sup> (+3), Cl <sup>-</sup> (-1)                                                                                        |
| <b>Rule 3</b> | Fluorine                   | <b>Always -1</b>             | Most electronegative element (No exceptions)                                                                                                                                    |
| <b>Rule 4</b> | Oxygen                     | <b>Usually -2</b>            | Peroxides (H <sub>2</sub> O <sub>2</sub> , Na <sub>2</sub> O <sub>2</sub> : -1); Superoxides (KO <sub>2</sub> : -1/2); OF <sub>2</sub> (+2); O <sub>2</sub> F <sub>2</sub> (+1) |
| <b>Rule 5</b> | Hydrogen                   | <b>Usually +1</b>            | Binary Metal Hydrides (LiH, NaH, CaH <sub>2</sub> : -1)                                                                                                                         |
| <b>Rule 6</b> | Halogens (Cl, Br, I)       | <b>Usually -1</b>            | Positive (+1, +3, +5, +7) when bonded to O or F                                                                                                                                 |
| <b>Rule 7</b> | Neutral Molecules / Ions   | <b>Σ O.N. = 0</b> (Molecule) | <b>Σ O.N. = Net Charge</b> (Polyatomic Ion)                                                                                                                                     |

**Stock Notation:**

• **Fe<sub>2</sub>O<sub>3</sub>** → Iron(III) oxide | **FeO** → Iron(II) oxide | **MnO<sub>2</sub>** → Manganese(IV) oxide | **AuCl<sub>3</sub>** → Gold(III) chloride

[Back to Interactive Index](#)

**4. STRUCTURAL PARADOXES & EXCEPTIONAL OXIDATION STATES**

Certain compounds yield fractional average oxidation numbers from chemical formulas because identical atoms occupy different bonding environments!

**1. Caro's Acid (H<sub>2</sub>SO<sub>5</sub> — Peroxomonosulphuric Acid)**

- *Formula Calculation:* 2(+1) + S + 5(-2) = 0 → S = +8 ( IMPOSSIBLE! Max valence of S is 6).
- *Structural Explanation:* Contains 1 Peroxy linkage (-O-O-, O.N. = -1). Structure: H-O-S(=O)<sub>2</sub>-O-O-H.
- *Correct Calculation:* 2(+1) [H] + S + 3(-2) [Oxo O] + 2(-1) [Peroxy O] = 0 → **S = +6** ( ).

**2. Chromium Peroxide (CrO<sub>5</sub> — Butterfly Structure)**

- *Formula Calculation:* Cr + 5(-2) = 0 → Cr = +10 ( IMPOSSIBLE!).

• *Structural Explanation:* Contains 4 peroxy oxygens (-1) and 1 oxo oxygen (-2).

• *Correct Calculation:*  $\text{Cr} + 1(-2) + 4(-1) = 0 \rightarrow \text{Cr} = +6$  ( ).

### 3. Fractional Oxidation State Examples

• **Fe<sub>3</sub>O<sub>4</sub> (Magnetite):** Mixture of FeO (+2) and Fe<sub>2</sub>O<sub>3</sub> (+3) → Average O.N. = **+8/3**.

• **Pb<sub>3</sub>O<sub>4</sub> (Red Lead):** Mixture of 2PbO (+2) and PbO<sub>2</sub> (+4) → Average O.N. = **+8/3**.

• **Br<sub>3</sub>O<sub>8</sub> (Tribromooctaoxide):** 2 terminal Br = +6; 1 central Br = +4 → Average O.N. = **+16/3**.

• **S<sub>4</sub>O<sub>6</sub><sup>2-</sup> (Tetrathionate Ion):** 2 terminal S = +5; 2 central S = 0 → Average O.N. = **+2.5**.

[Back to Interactive Index](#)

## 5. TYPES OF REDOX REACTIONS & DISPROPORTIONATION

| Type of Redox Reaction | General Equation                                 | Classic NCERT Example                                                                        | Key Feature                                                |
|------------------------|--------------------------------------------------|----------------------------------------------------------------------------------------------|------------------------------------------------------------|
| Combination            | $A + B \rightarrow C$                            | $\text{C(s)} + \text{O}_2\text{(g)} \rightarrow \text{CO}_2\text{(g)}$                       | At least one reactant must be in elemental form            |
| Decomposition          | $C \rightarrow A + B$                            | $2\text{H}_2\text{O(l)} \rightarrow 2\text{H}_2\text{(g)} + \text{O}_2\text{(g)}$            | At least one product must be in elemental form             |
| Metal Displacement     | $X + \text{YZ} \rightarrow \text{XZ} + \text{Y}$ | $\text{CuSO}_4 + \text{Zn} \rightarrow \text{ZnSO}_4 + \text{Cu}$                            | More reactive metal displaces less reactive metal          |
| Non-Metal Displacement | $X + \text{YZ} \rightarrow \text{XZ} + \text{Y}$ | $2\text{Na} + 2\text{H}_2\text{O} \rightarrow 2\text{NaOH} + \text{H}_2$                     | Active metal displaces H <sub>2</sub> gas from water/ acid |
| Disproportionation     | $A \rightarrow B + C$                            | $2\text{H}_2\text{O}_2\text{(aq)} \rightarrow 2\text{H}_2\text{O(l)} + \text{O}_2\text{(g)}$ | <b>SAME element is oxidised and reduced simultaneously</b> |

#### EDUVISTA EXAM ALERT (FLUORINE DISPROPORTIONATION):

Fluorine (F<sub>2</sub>) **CANNOT** undergo disproportionation because it is the most electronegative element and exhibits only 0 and -1 oxidation states!

[Back to Interactive Index](#)

## 6. BALANCING REDOX WORKBENCH

### STEP-BY-STEP BALANCING WORKBENCH: ACIDIC MEDIUM

**Target Reaction:**  $\text{MnO}_4^-\text{(aq)} + \text{Fe}^{2+}\text{(aq)} \rightarrow \text{Mn}^{2+}\text{(aq)} + \text{Fe}^{3+}\text{(aq)}$

**Step 1: Split into Half-Reactions**Oxidation:  $\text{Fe}^{2+} \rightarrow \text{Fe}^{3+}$  | Reduction:  $\text{MnO}_4^- \rightarrow \text{Mn}^{2+}$ **Step 2: Balance Oxygen with  $\text{H}_2\text{O}$** Reduction:  $\text{MnO}_4^- \rightarrow \text{Mn}^{2+} + 4\text{H}_2\text{O}$ **Step 3: Balance Hydrogen with  $\text{H}^+$** Reduction:  $\text{MnO}_4^- + 8\text{H}^+ \rightarrow \text{Mn}^{2+} + 4\text{H}_2\text{O}$ **Step 4: Balance Net Charge with  $\text{e}^-$** Oxidation:  $\text{Fe}^{2+} \rightarrow \text{Fe}^{3+} + \text{e}^- (\times 5)$ Reduction:  $\text{MnO}_4^- + 8\text{H}^+ + 5\text{e}^- \rightarrow \text{Mn}^{2+} + 4\text{H}_2\text{O} (\times 1)$ **Step 5: Add Half-Reactions & Cancel  $\text{e}^-$** **Final Balanced Equation:**  $\text{MnO}_4^- + 5\text{Fe}^{2+} + 8\text{H}^+ \rightarrow \text{Mn}^{2+} + 5\text{Fe}^{3+} + 4\text{H}_2\text{O}$ [Back to Interactive Index](#)

## 7. REDOX TITRATIONS & INDICATORS

**REDOX TITRATION INDICATOR MECHANISMS**

- |                          |                                                                                                                                      |
|--------------------------|--------------------------------------------------------------------------------------------------------------------------------------|
| 1. SELF-INDICATOR        | : $\text{KMnO}_4$ (Purple $\text{MnO}_4^- \rightarrow$ Colourless $\text{Mn}^{2+}$ ; Pink endpoint)                                  |
| 2. INTERNAL INDICATOR    | : $\text{K}_2\text{Cr}_2\text{O}_7$ (Diphenylamine indicator $\rightarrow$ Intense Blue-Violet)                                      |
| 3. IODOMETRIC TITRATION: | $\text{I}_2 + 2\text{S}_2\text{O}_3^{2-} \rightarrow 2\text{I}^- + \text{S}_4\text{O}_6^{2-}$ (Starch $\rightarrow$ Blue-Colourless) |

[Back to Interactive Index](#)

## 8. ELECTROCHEMICAL CELLS & DANIELL CELL

A **Galvanic (Electrochemical) Cell** converts chemical energy of a spontaneous redox reaction into electrical energy.

**EDUVISTA MEMORY TRICK FOR ANODE: "LOAN"**

- **L** — Left side
- **O** — Oxidation occurs
- **A** — Anode electrode
- **N** — Negative polarity

**Functions of Salt Bridge:**

1. Completes electrical circuit without allowing mixing of half-cell solutions.
2. Maintains electrical neutrality in both half-cell compartments via ion migration ( $\text{K}^+$ ,  $\text{Cl}^-$ ,  $\text{NO}_3^-$ ).

[Back to Interactive Index](#)

## 9. STANDARD ELECTRODE POTENTIAL & ELECTROCHEMICAL SERIES

• **Cell EMF Formula:**

$$E^{\circ}_{\text{cell}} = E^{\circ}_{\text{cathode}} - E^{\circ}_{\text{anode}} = E^{\circ}_{\text{Right}} - E^{\circ}_{\text{Left}}$$

• **Spontaneity Criterion:** Reaction is spontaneous if  $E^{\circ}_{\text{cell}} > 0$ .

| ELECTROCHEMICAL SERIES HIGHLIGHTS                                                             |                               |                           |
|-----------------------------------------------------------------------------------------------|-------------------------------|---------------------------|
| $\text{F}_2(\text{g}) + 2\text{e}^- \rightarrow 2\text{F}^-$                                  | $E^{\circ} = +2.87 \text{ V}$ | Strongest Oxidising Agent |
| $\text{MnO}_4^- + 8\text{H}^+ + 5\text{e}^- \rightarrow \text{Mn}^{2+} + 4\text{H}_2\text{O}$ | $E^{\circ} = +1.51 \text{ V}$ | Strong Oxidising Agent    |
| $2\text{H}^+ + 2\text{e}^- \rightarrow \text{H}_2(\text{g})$                                  | $E^{\circ} = 0.00 \text{ V}$  | REFERENCE (SHE)           |
| $\text{Zn}^{2+} + 2\text{e}^- \rightarrow \text{Zn}(\text{s})$                                | $E^{\circ} = -0.76 \text{ V}$ | Reducing Agent            |
| $\text{Li}^+ + \text{e}^- \rightarrow \text{Li}(\text{s})$                                    | $E^{\circ} = -3.05 \text{ V}$ | Strongest Reducing Agent  |

[Back to Interactive Index](#)

## 10. CBSE BOARD EXAM PREVIOUS YEAR QUESTIONS (SOLVED)

**CBSE 2024 [1 MARK]**

**Q1.** Identify the oxidising agent in:  $\text{H}_2\text{S}(\text{g}) + \text{Cl}_2(\text{g}) \rightarrow 2\text{HCl}(\text{g}) + \text{S}(\text{s})$

[Click to View Solution & Marking Scheme](#)

**Solution:** Chlorine ( $\text{Cl}_2$ ) is reduced from oxidation state 0 (in  $\text{Cl}_2$ ) to -1 (in  $\text{HCl}$ ). Therefore,  **$\text{Cl}_2$  is the oxidising agent.**

**CBSE 2023 [2 MARKS]**

**Q2.** Calculate the oxidation number of the underlined element: (a)  $\text{K}_3[\text{Fe}(\text{CN})_6]$  (b)  $\text{CrO}_5$

[Click to View Solution & Marking Scheme](#)

(a)  $\text{K}_3[\text{Fe}(\text{CN})_6]$ :  $3(+1) + x + 6(-1) = 0 \Rightarrow 3 + x - 6 = 0 \Rightarrow x = +3$ .

(b)  $\text{CrO}_5$ : Structure has 4 peroxy oxygens (-1) and 1 oxo oxygen (-2)  $\Rightarrow x + 1(-2) + 4(-1) = 0 \Rightarrow x = +6$ .

**CBSE 2023 [3 MARKS]**

**Q3.** Balance in acidic medium:  $\text{MnO}_4^-(\text{aq}) + \text{Fe}^{2+}(\text{aq}) \rightarrow \text{Mn}^{2+}(\text{aq}) + \text{Fe}^{3+}(\text{aq})$

[Click to View Solution & Marking Scheme](#)

Oxidation:  $\text{Fe}^{2+} \rightarrow \text{Fe}^{3+} + \text{e}^- (\times 5)$

Reduction:  $\text{MnO}_4^- + 8\text{H}^+ + 5\text{e}^- \rightarrow \text{Mn}^{2+} + 4\text{H}_2\text{O} (\times 1)$

**Final Balanced Equation:**  $\text{MnO}_4^- + 5\text{Fe}^{2+} + 8\text{H}^+ \rightarrow \text{Mn}^{2+} + 5\text{Fe}^{3+} + 4\text{H}_2\text{O}$

CBSE 2022 [3 MARKS]

**Q4. Balance in basic medium:**  $\text{MnO}_4^-(\text{aq}) + \text{I}^-(\text{aq}) \rightarrow \text{MnO}_2(\text{s}) + \text{I}_2(\text{s})$

[Click to View Solution & Marking Scheme](#)

Oxidation:  $2\text{I}^- \rightarrow \text{I}_2 + 2\text{e}^-$  ( $\times 3$ )

Reduction:  $\text{MnO}_4^- + 4\text{H}^+ + 3\text{e}^- \rightarrow \text{MnO}_2 + 2\text{H}_2\text{O}$  ( $\times 2$ )

Add 8  $\text{OH}^-$  to both sides & simplify:  $2\text{MnO}_4^- + 6\text{I}^- + 4\text{H}_2\text{O} \rightarrow 2\text{MnO}_2 + 3\text{I}_2 + 8\text{OH}^-$

CBSE 2020 [5 MARKS]

**Q5. Justify why**  $2\text{H}_2\text{O}_2(\text{aq}) \rightarrow 2\text{H}_2\text{O}(\text{l}) + \text{O}_2(\text{g})$  **is a disproportionation reaction.**

[Click to View Solution & Marking Scheme](#)

- Oxygen in  $\text{H}_2\text{O}_2$  has oxidation number **-1**.
- In  $\text{H}_2\text{O}$ , oxygen decreases to **-2** (Reduction).
- In  $\text{O}_2$ , oxygen increases to **0** (Oxidation).

Since the **same element (Oxygen)** is **simultaneously oxidised and reduced**, this is a disproportionation reaction.

[Back to Interactive Index](#)

## 11. JEE MAIN & NEET COMPETITIVE PYQS (SOLVED)

JEE MAIN 2023

**Q1. The oxidation state of central sulfur atom in  $\text{S}_4\text{O}_6^{2-}$  and Caro's acid ( $\text{H}_2\text{SO}_5$ ) respectively are:**

[Click to View Solution & Analysis](#)

**Correct Answer: (B) or (C)** [Terminal S = +5, Central S = 0 in  $\text{S}_4\text{O}_6^{2-}$ ; S = +6 in  $\text{H}_2\text{SO}_5$ ].

In  $\text{S}_4\text{O}_6^{2-}$ , the structure is  $^-\text{O}_3\text{S}-\text{S}-\text{S}-\text{SO}_3^-$ . Terminal sulfur atoms are +5, central are 0. Caro's acid ( $\text{H}_2\text{SO}_5$ ) has 1 peroxy linkage  $\rightarrow \text{S} = +6$ .

NEET 2022

**Q2. Which halogen does NOT undergo disproportionation in alkaline medium?**

[Click to View Solution & Analysis](#)

**Correct Answer: (D) F<sub>2</sub>**

Fluorine is the most electronegative element and exhibits only 0 and -1 oxidation states. It cannot exhibit positive oxidation states needed for disproportionation.

**JEE MAIN 2022**

**Q3. For Zn(s) | Zn<sup>2+</sup>(aq) || Cu<sup>2+</sup>(aq) | Cu(s), given E°(Zn<sup>2+</sup>/Zn) = -0.76 V and E°(Cu<sup>2+</sup>/Cu) = +0.34 V. Calculate E°cell.**

[Click to View Solution & Analysis](#)

E°cell = E°cathode - E°anode = (+0.34 V) - (-0.76 V) = **+1.10 V** (Spontaneous).

**NEET 2021**

**Q4. Volume of 0.02 M KMnO<sub>4</sub> required to oxidise 20 mL of 0.1 M FeSO<sub>4</sub> in acidic medium?**

[Click to View Solution & Analysis](#)

$M_1 \times V_1 \times n_1 = M_2 \times V_2 \times n_2 \Rightarrow 0.02 \times V_1 \times 5 = 0.1 \times 20 \times 1 \Rightarrow 0.1 \times V_1 = 2 \Rightarrow V_1 = 20 \text{ mL}.$

[Back to Interactive Index](#)

## 12. INTERACTIVE SELF-TEST PRACTICE QUIZ

### TEST YOUR KNOWLEDGE (5-MINUTE CHECKPOINT)

**Q1: What is the oxidation number of Cr in CrO<sub>5</sub>?**

[Reveal Answer](#)

**Answer: +6** (Contains 4 peroxy oxygens of -1 and 1 oxo oxygen of -2).

**Q2: Name the indicator used in Iodometric Titrations.**

[Reveal Answer](#)

**Answer: Starch Indicator** (Forms deep blue complex with I<sub>2</sub>).

**Q3: What does the LOAN rule stand for in Galvanic cells?**

**Reveal Answer**

**Answer: Left, Oxidation, Anode, Negative.**

[Back to Interactive Index](#)

## EDUVISTA ACADEMIC PROGRAMS — ADMISSIONS OPEN!

Personalised Learning • Smart Notes • Concept Clarity • Guaranteed Results

**Direct Helpline / WhatsApp: 8920696949**

**Website & Portal: Eduvista Smart Notes Learning System**